

RAGNAR-OLAF BUCHWEITZ AND SHIPING LIU

ABSTRACT. We show that the dimension of the mesh algebra of a finite Auslander-Reiten quiver over a field is a purely combinatorial invariant that does not depend on the ground field. Moreover, a combinatorial algorithm for computing this dimension is given along our proof of this result.

Translation quivers appear naturally in the representation theory of finite dimensional algebras; see, for example, [3]. A translation quiver defines a mesh algebra over any field. A natural question arises as to whether or not the dimension of the mesh algebra depends on the field. The purpose of this note is to show that the dimension of the mesh algebra of a finite Auslander-Reiten quiver over a field is a purely combinatorial invariant of this quiver. Indeed, our proof yields a combinatorial algorithm for computing this dimension. As a further application, one may use then semicontinuity of Hochschild cohomology of algebras as in [6] to conclude that a finite Auslander-Reiten quiver contains no oriented cycle if its mesh algebra over some field admits no outer derivation.

1. *Simply connected translation quivers*

Throughout this note, Γ denotes a translation quiver with translation τ ; see [3, (1.1)]. We assume that Γ contains neither loops nor multiple arrows but that each non-projective vertex is the end-point of at least one arrow. One defines the *orbit graph* $\mathcal{O}(\Gamma)$ of Γ as follows: the τ -orbit of a vertex a is the set $o(x)$ of vertices of the form $\tau^n(x)$ with $n \in \mathbb{Z}$; the vertices of $\mathcal{O}(\Gamma)$ are the τ -orbits of Γ , and there exists an edge $o(x) - o(y)$ in $\mathcal{O}(\Gamma)$ if Γ contains an arrow $a \rightarrow b$ or $b \rightarrow a$ with $a \in o(x)$ and $b \in o(y)$. If Γ contains no oriented cycle, then $\mathcal{O}(\Gamma)$ is the graph G_Γ defined in [3, (4.2)]. Now Γ is called *simply connected* if Γ contains no oriented cycle and $\mathcal{O}(\Gamma)$ is a tree. By [3, (1.6), (4.1), (4.2)], this definition is equivalent to that given in [3, (1.6)].

Let $p : x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_r$ be a path in Γ . Then p induces a walk $w(p) : o(x_0) - o(x_1) - \cdots - o(x_r)$ in $\mathcal{O}(\Gamma)$, and $w(p)$ in turn determines a unique reduced walk $w_{\text{red}}(p)$. Recall that the path p is called *sectional* if

$x_{i-1} \neq \tau x_{i+1}$ for each $0 < i < r$. Moreover, for a vertex $x \in \Gamma$, we denote by x^+ and by x^- the set of immediate successors and that of immediate predecessors of x , respectively.

We recall that a sequence

$$a_0 \xrightarrow{e_1} a_1 \cdots \xrightarrow{e_r} a_r$$

of edges of a graph such that $e_i \neq e_{i+1}$ for all $1 \leq i < r$ is called a *reduced walk*. A vertex of a graph is considered as a *trivial* reduced walk.

1.1. LEMMA. *Assume that Γ contains no oriented cycle.*

(1) *Let x be a vertex of Γ . If $x_1, x_2 \in x^-$ or $x_1, x_2 \in x^+$, then $o(x_1) = o(x_2)$ if and only if $x_1 = x_2$.*

(2) *If p is a sectional path in Γ , then $w(p)$ is a reduced walk in $\mathcal{O}(\Gamma)$.*

Proof. (1) It suffices to consider the case where $x_1, x_2 \in x^-$. Suppose that $o(x_1) = o(x_2)$. We may assume that $x_2 = \tau^r x_1$ for some $r \geq 0$. If $r > 0$, then $x \rightarrow \tau^{r-1} x_1 \rightarrow \cdots \rightarrow x_1 \rightarrow x$ would be an oriented cycle. Thus $r = 0$, that is $x_1 = x_2$.

(2) Let $p : x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_r$ be a sectional path in Γ . If $w(p)$ is not reduced, then $o(x_{i-1}) = o(x_{i+1})$ for some $0 < i < r$. However, this means that $x_{i-1} = \tau^m x_{i+1}$ for some integer m . As Γ contains no oriented cycle, m is necessarily positive. In particular, τx_{i+1} exists, and then by part (1), $x_{i-1} = \tau x_{i+1}$ since $x_{i-1}, \tau x_{i+1} \in x_i^-$. This contradicts the hypothesis that p is sectional. The proof of the lemma is completed.

1.2. LEMMA. *If Γ is simply connected, then a sectional path in Γ is the only path between its endpoints.*

Proof. Assume that Γ is simply connected. We shall prove the lemma by induction on the length of a sectional path. The lemma is trivially true if the length is zero, as Γ contains no oriented cycle. Assume that $r > 0$ and the lemma holds for sectional paths of length less than r . Let

$$p : x = y_0 \rightarrow y_1 \rightarrow \cdots \rightarrow y_{r-1} \rightarrow y_r = y$$

be a sectional path. Then the path u obtained from p by dropping the initial arrow is the unique path from y_1 to y . Note that u does not start with $y_1 \rightarrow \tau^- x$ as p is sectional. Now let

$$q : x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_{s-1} \rightarrow x_s = y$$

be another path in Γ from x to y . We claim first that $o(x) \neq o(x_t)$ for $0 < t \leq s$. Otherwise, $x_t = \tau^{-m} x$ for some such t and some integer m . Using again the

fact that Γ contains no oriented cycle, we have $m > 0$ and consequently a path from τ^-x to $\tau^{-m}x$. Further the initial arrow $x \rightarrow y_1$ gives rise to an arrow $y_1 \rightarrow \tau^-x$. Thus we obtain a path

$$y_1 \rightarrow \tau^-x \rightarrow \cdots \rightarrow \tau^{-m}x = x_t \rightarrow x_{t+1} \rightarrow \cdots \rightarrow x_{s-1} \rightarrow y,$$

which is different from u , a contradiction.

Thus, $o(x) \neq o(x_t)$ for $0 < t \leq s$, as claimed. It follows that $o(x) - o(x_1)$ is the initial edge of the reduced walk $w_{\text{red}}(q)$ from $o(x)$ to $o(y)$ in $\mathcal{O}(\Gamma)$. Now $w(p)$ is a reduced walk from $o(x)$ to $o(y)$ whose initial edge is $o(x) - o(y_1)$. As $\mathcal{O}(\Gamma)$ is a tree, the edge $o(x) - o(x_1)$ coincides with the edge $o(x) - o(y_1)$. In particular, $o(x_1) = o(y_1)$. Thus $x_1 = y_1$, by Lemma 1.1.(1). Since Γ contains no multiple arrow, the initial arrow $x \rightarrow x_1$ of q is the same as that of p . Hence $p = q$ by the inductive hypothesis. This completes the proof of the lemma.

For a vertex x of Γ , we define the *forward cone* Γ_x of x to be the set of all successors of x in Γ .

1.3. LEMMA. *Assume that Γ is simply connected. Let x be a vertex of Γ and assume that*

$$p : x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_{m-1} \rightarrow x_m$$

is a non-trivial sectional path in Γ with $x_0 \in \Gamma_x$ and x_1 projective. For each $1 \leq i \leq m$, the vertex x_{i-1} is then the only immediate predecessor of x_i that belongs to Γ_x .

Proof. As $x_0 \in \Gamma_x$, there exists a path

$$q : x = y_0 \rightarrow y_1 \rightarrow \cdots \rightarrow y_{r-1} \rightarrow y_r = x_0 \rightarrow x_1$$

in Γ . Since x_1 is projective and Γ contains no oriented cycle, $o(y_i) \neq o(x_1)$ for all $0 \leq i \leq r$. Therefore, the associated reduced walk $w_{\text{red}}(q)$ from $o(x)$ to $o(x_1)$ in $\mathcal{O}(\Gamma)$ ends with the edge $o(x_0) - o(x_1)$. If

$$u : x = a_0 \rightarrow a_1 \rightarrow \cdots \rightarrow a_{s-1} \rightarrow a_s = x_1$$

is a second such path, by the same argument, the reduced walk $w_{\text{red}}(u)$ from $o(x)$ to $o(x_1)$ ends with the edge $o(a_{s-1}) - o(x_1)$. This reduced walk coincides with the reduced walk $w_{\text{red}}(q)$, as $\mathcal{O}(\Gamma)$ is a tree. Consequently, $o(a_{s-1}) = o(x_0)$, and then, by Lemma 1.1.(1), $a_{s-1} = x_0$. This shows that the lemma holds for $m = 1$. Assume now that $m > 1$ and that the lemma holds for $m - 1$. By Lemma 1.1.(2),

$$w(p) : o(x_0) - o(x_1) - \cdots - o(x_{m-1}) - o(x_m)$$

is a reduced walk in $\mathcal{O}(\Gamma)$. Combining $w(p)$ with the reduced walk $w_{\text{red}}(q)$ results in a reduced walk

$$w : o(x) \text{ --- } \cdots \text{ --- } o(x_0) \text{ --- } o(x_1) \text{ --- } \cdots \text{ --- } o(x_{m-1}) \text{ --- } o(x_m).$$

Let $v : x = z_0 \rightarrow z_1 \rightarrow \cdots \rightarrow z_{t-1} \rightarrow z_t = x_m$ be a path in Γ . If $o(x_m) = o(z_i)$ for some $0 \leq i < t$, then $z_i = \tau^n x_m$. Now $n > 0$ as Γ contains no oriented cycle. This implies that τx_m is in Γ_x , and then τx_m and x_{m-2} were different immediate predecessors of x_{m-1} in Γ_x , contrary to the inductive hypothesis. Thus, $o(x_m) \neq o(z_i)$ for each $0 \leq i < t$, and so u induces a reduced walk from $o(x)$ to $o(x_m)$ that ends with the edge $o(z_{t-1}) \text{ --- } o(x_m)$. Again $o(z_{s-1}) = o(x_{m-1})$ since $\mathcal{O}(\Gamma)$ is a tree, and hence $z_{s-1} = x_{m-1}$, by Lemma 1.1. This completes the induction, thus, the proof of the lemma.

1.4. LEMMA. *Assume that Γ is simply connected. If x, y are vertices of Γ , then all paths from x to y in Γ are of the same length.*

Proof. We fix a vertex $x \in \Gamma$. Let $x = x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_r = y$ be a path in Γ . We shall use induction on i to prove that all paths from x to x_i are of length i . This is trivially true for $i = 0$ since Γ contains no oriented cycle. Assume that $1 \leq i \leq r$ and all paths from x to x_{i-1} are of length $i - 1$. Assume further that

$$q : x = y_0 \rightarrow y_1 \rightarrow \cdots \rightarrow y_s = x_i$$

is another path. Then $s > 0$. Let t with $0 \leq t < s$ be the smallest integer such that the path $y_t \rightarrow \cdots \rightarrow y_s$ is sectional. If $t = 0$, then $s = i$ by Lemma 1.2. Assume that $t > 0$. Then $y_{t-1} = \tau y_{t+1}$ by the minimality of t . If some y_j with $t + 1 < j \leq s$ is projective, then $y_{s-1} = x_{i-1}$ by Lemma 1.3. Therefore, $s - 1 = i - 1$ by the inductive hypothesis. Otherwise, Γ contains a path

$$x \rightarrow y_1 \rightarrow \cdots \rightarrow y_{t-1} = \tau y_{t+1} \rightarrow \cdots \rightarrow \tau y_s \rightarrow x_{i-1}.$$

By the inductive hypothesis, $i - 1 = (t - 1) + (s - t)$, whence $s = i$. This completes the induction, and hence the proof of the lemma.

2. Mesh algebras

The objective of this section is to prove our main result. For this purpose, we need to recall some basic notions and results on linear categories from [3]. Throughout this section, k denotes an algebraically closed field and Λ a locally bounded k -category. There exists a unique locally finite quiver Q_Λ , called the *ordinary quiver* of Λ , such that $\Lambda \cong kQ_\Lambda/I_\Lambda$, where kQ_Λ is the path category of Q_Λ over k and I_Λ is an ideal contained in the ideal of kQ_Λ generated by the

paths of length two. The *radical* $\mathfrak{R}\Lambda$ of Λ is the ideal generated by the non-invertible morphisms. Set $\mathfrak{R}^0\Lambda = \Lambda$ and define $\mathfrak{R}^n\Lambda$ for $n \geq 1$ to be the ideal generated by the composites of n morphisms from $\mathfrak{R}\Lambda$. Denote by $\text{mod}\Lambda$ the category of finite dimensional right Λ -modules and by $\text{ind}\Lambda$ the full subcategory of $\text{mod}\Lambda$ generated by a chosen complete set of representatives of isomorphism classes of indecomposable modules in $\text{mod}\Lambda$.

We shall denote by $k(\Gamma)$ the *mesh category* of Γ over k ; see [3, (2.5)]. Note that for each $n \geq 0$, the ideal $\mathfrak{R}^n k(\Gamma)$ is generated by the classes of paths of length n in Γ modulo the mesh ideal. If Γ is finite, then $\bigoplus_{x,y \in \Gamma} k(\Gamma)(x,y)$ is a k -algebra, called the *mesh algebra* of Γ over k and denoted again by $k(\Gamma)$. Note that we can define in the same way the mesh algebra $R(\Gamma)$ of Γ over any commutative ring R .

Assume now that Λ is *locally representation-finite*, that is, every object x of Λ lies in the support of only finitely many modules in $\text{ind}\Lambda$, or equivalently, $\text{ind}\Lambda$ is locally bounded. The Auslander-Reiten quiver Γ_Λ of Λ is a translation quiver with respect to the Auslander-Reiten translation $\tau_\Lambda = DTr$, the dual of the transpose. Recall that Λ is called *standard*, [3, (5.1)], if $\text{ind}\Lambda \cong k(\Gamma_\Lambda)$, and *simply connected*, [3, (6.1)], if Γ_Λ is a simply connected translation quiver. It follows from [4, (3.1)] that standardness in this sense is equivalent to that defined in [2, (1.11)]. If Γ_Λ contains no oriented cycle, then the ordinary quiver Q_Λ of Λ contains no loop as $\text{ind}\Lambda$ is locally bounded, and consequently, Λ is standard [2, (9.6)]. In particular, Λ is standard if it is simply connected.

We say that Γ is an *Auslander-Reiten quiver over k* if $\Gamma \cong \Gamma_\Lambda$ for some locally representation-finite k -category Λ . In this case, $k(\Gamma)$ is a locally bounded k -category. It follows from Brenner's result [5, (5.1)] that if Γ is an Auslander-Reiten quiver over k , then it is an Auslander-Reiten quiver over any algebraically closed field. In the sequel, we shall simply say that Γ is an Auslander-Reiten quiver with no reference to a ground field.

LEMMA 2.1. *Let Γ be a simply connected Auslander-Reiten quiver, and let k, l be algebraically closed fields. For any pair x, y of vertices of Γ and any integer $n \geq 0$, one has then*

$$\dim_k \mathfrak{R}^n k(\Gamma)(x, y) = \dim_l \mathfrak{R}^n l(\Gamma)(x, y).$$

Proof. Let Λ and Σ be a locally representation-finite k -category and l -category, respectively, such that $\Gamma \cong \Gamma_\Lambda \cong \Gamma_\Sigma$. Then Λ and Σ are simply connected, in particular, Λ and Σ are standard. Thus $k(\Gamma) \cong k(\Gamma_\Lambda) \cong \text{ind}\Lambda$, and similarly $l(\Gamma) \cong \text{ind}\Sigma$.

We fix a vertex x of Γ . If $y \notin \Gamma_x$, then $\dim_k k(\Gamma)(x, y) = 0 = \dim_l l(\Gamma)(x, y)$. Assume now that $y \in \Gamma_x$. By Lemma 1.4, all paths from x to y have the same length, which we denote by $d(x, y)$.

If $d(x, y) = 0$, then $\dim_k k(\Gamma)(x, y) = 1 = \dim_l l(\Gamma)(x, y)$. Now assume that $d(x, y) = r > 0$ and $\dim_k k(\Gamma)(x, z) = \dim_l l(\Gamma)(x, z)$ for all $z \in \Gamma_x$ with $d(x, z) < r$. Assume that $y^- = \{x_1, \dots, x_s\}$. For each $1 \leq i \leq s$, either $x_i \notin \Gamma_x$, or $x_i \in \Gamma_x$ and then $d(x, x_i) = r - 1$. It follows thus from the inductive hypothesis that $\dim_k k(\Gamma)(x, x_i) = \dim_l l(\Gamma)(x, x_i)$ for all $1 \leq i \leq s$.

Let $\phi : \Gamma \rightarrow \Gamma_\Lambda$ be an isomorphism that exists by our assumption. Set $M = \phi(x)$, $N = \phi(y)$, and $M_i = \phi(x_i)$ for $1 \leq i \leq s$. If y is not a projective vertex, then N is not a projective module. There exists thus an almost split sequence

$$(*) \quad 0 \rightarrow \tau_\Lambda N \rightarrow M_1 \oplus \dots \oplus M_s \rightarrow N \rightarrow 0$$

in $\text{mod } \Lambda$. Note that $M \not\cong N$, since Γ_Λ contains no oriented cycle and $r > 0$. Accordingly, applying $\text{Hom}_\Lambda(M, -)$ to the above almost split sequence yields an exact sequence

$$0 \rightarrow \text{Hom}_\Lambda(M, \tau_\Lambda N) \rightarrow \bigoplus_{i=1}^s \text{Hom}_\Lambda(M, M_i) \rightarrow \text{Hom}_\Lambda(M, N) \rightarrow 0,$$

and counting dimensions gives rise to the following equalities:

$$\begin{aligned} \dim_k k(\Gamma)(x, y) &= \dim_k \text{Hom}_\Lambda(M, N) \\ &= \sum_{i=1}^s \dim_k \text{Hom}_\Lambda(M, M_i) - \dim_k \text{Hom}_\Lambda(M, \tau_\Lambda N) \\ &= \sum_{i=1}^s \dim_k k(\Gamma)(x, x_i) - \dim_k k(\Gamma)(x, \tau y). \end{aligned}$$

If y is a projective vertex, then N is a projective module, and so

$$\begin{aligned} \dim_k k(\Gamma)(x, y) &= \dim_k \text{Hom}_\Lambda(M, N) \\ &= \sum_{i=1}^s \dim_k \text{Hom}_\Lambda(M, M_i) \\ &= \sum_{i=1}^s \dim_k k(\Gamma)(x, x_i). \end{aligned}$$

Similarly, $\dim_l l(\Gamma)(x, y) = \sum_{i=1}^s \dim_l l(\Gamma)(x, x_i) - \dim_l l(\Gamma)(x, \tau y)$, if y is non-projective; and otherwise, $\dim_l l(\Gamma)(x, y) = \sum_{i=1}^s \dim_l l(\Gamma)(x, x_i)$. In case y is non-projective, either $\tau y \notin \Gamma_x$ or $\tau y \in \Gamma_x$ with $d(x, \tau y) = r - 2$. Thus $\dim_k k(\Gamma)(x, \tau y) = \dim_l l(\Gamma)(x, \tau y)$ by the inductive hypothesis. Therefore, $\dim_k k(\Gamma)(x, y) = \dim_l l(\Gamma)(x, y)$ in either case.

Finally, let us fix an integer $n \geq 1$. If Γ contains no path from x to y of length $\geq n$, then $\mathfrak{R}^n k(\Gamma)(x, y) = 0 = \mathfrak{R}^n l(\Gamma)(x, y)$. Otherwise, by Lemma 1.4, all paths from x to y are of length $\geq n$. Hence $\mathfrak{R}^n k(\Gamma)(x, y) = k(\Gamma)(x, y)$ and $\mathfrak{R}^n l(\Gamma)(x, y) = l(\Gamma)(x, y)$. Thus $\dim_k \mathfrak{R}^n k(\Gamma)(x, y) = \dim_l \mathfrak{R}^n l(\Gamma)(x, y)$. This completes the proof of the lemma.

Note that if Γ is a finite Auslander-Reiten quiver, then $k(\Gamma)$ is finite dimensional over k . In order to show that the mesh algebra $F(\Gamma)$ of Γ over any other

field F is of dimension $\dim_k k(\Gamma)$, recall that a k -linear functor $\pi : \tilde{\Lambda} \rightarrow \Lambda$ of locally bounded k -categories is a *covering functor*, [3, (3.1)], if for all objects x, y of Λ , the fibre $\pi^{-}(x)$ is non-empty, and for each $\tilde{x} \in \pi^{-}(x)$, the map π induces k -isomorphisms

$$\bigoplus_{\tilde{y} \in \pi^{-}(y)} \tilde{\Lambda}(\tilde{x}, \tilde{y}) \xrightarrow{\cong} \Lambda(x, y) \quad \text{and} \quad \bigoplus_{\tilde{y} \in \pi^{-}(y)} \tilde{\Lambda}(\tilde{y}, \tilde{x}) \xrightarrow{\cong} \Lambda(y, x).$$

It is easy to see that π induces, for each $n \geq 1$, a k -isomorphism

$$\bigoplus_{\tilde{y} \in \pi^{-}(y)} \mathfrak{R}^n \tilde{\Lambda}(\tilde{x}, \tilde{y}) \xrightarrow{\cong} \mathfrak{R}^n \Lambda(x, y).$$

2.2. THEOREM. *Let Γ be a finite Auslander-Reiten quiver. For each $n \geq 0$, there exists $d_n \geq 0$ such that $\dim_F \mathfrak{R}^n F(\Gamma) = d_n$ for any field F .*

Proof. We fix an algebraically closed field k , and let F be an arbitrary field with algebraic closure l . Clearly, $\mathfrak{R}^n l(\Gamma) = \mathfrak{R}^n F(\Gamma) \otimes_F l$ for each $n \geq 0$, whence we need only to show that $\dim_k \mathfrak{R}^n k(\Gamma) = \dim_l \mathfrak{R}^n l(\Gamma)$ for each $n \geq 0$. Furthermore, it suffices to show that $\dim_k \mathfrak{R}^n k(\Gamma)(x, y) = \dim_l \mathfrak{R}^n l(\Gamma)(x, y)$ for each pair x, y of vertices in Γ and each $n \geq 0$. Let $\pi : \tilde{\Gamma} \rightarrow \Gamma$ be the universal covering of Γ ; see [3, (1.3)]. Then π induces covering functors $k(\tilde{\Gamma}) \rightarrow k(\Gamma)$ and $l(\tilde{\Gamma}) \rightarrow l(\Gamma)$; see [3, (3.1)].

Let now Λ be a representation-finite k -category such that $\Gamma \cong \Gamma_\Lambda$. Let $\tilde{\Lambda}$ be the full subcategory of $k(\tilde{\Gamma})$ generated by the projective vertices. According to [3, (2.4)], $\tilde{\Lambda}$ is a locally representation-finite k -category such that $\Gamma_{\tilde{\Lambda}} \cong \tilde{\Gamma}$. We fix a vertex $\tilde{x} \in \pi^{-}(x)$ and an integer $n \geq 0$. Then

$$\mathfrak{R}^n k(\Gamma)(x, y) \cong \bigoplus_{\tilde{y} \in \pi^{-}(y)} \mathfrak{R}^n k(\tilde{\Gamma})(\tilde{x}, \tilde{y}).$$

Therefore, $\dim_k \mathfrak{R}^n k(\Gamma)(x, y) = \sum_{\tilde{y} \in \pi^{-}(y)} \dim_k \mathfrak{R}^n k(\tilde{\Gamma})(\tilde{x}, \tilde{y})$. Similarly, one can show that

$$\dim_l \mathfrak{R}^n l(\Gamma)(x, y) = \sum_{\tilde{y} \in \pi^{-}(y)} \dim_l \mathfrak{R}^n l(\tilde{\Gamma})(\tilde{x}, \tilde{y}).$$

Now by Lemma 2.1, $\dim_k \mathfrak{R}^n k(\tilde{\Gamma})(\tilde{x}, \tilde{y}) = \dim_l \mathfrak{R}^n l(\tilde{\Gamma})(\tilde{x}, \tilde{y})$, as $\tilde{\Gamma}$ is simply connected. This completes the proof of the theorem.

REMARK. It seems plausible that the above result should hold true for any translation quiver Γ with $k(\Gamma)$ being finite dimensional. However, the analogue to the exact sequence (*) in the proof of Lemma 2.1 fails for a general translation quiver.

We shall now show that the mesh algebra of a finite Auslander-Reiten quiver over a commutative ring is free of finite rank. For this purpose, we need the following basic result, in which all tensor products are taken over \mathbb{Z} .

2.3. LEMMA. Let A be a \mathbb{Z} -algebra, and let I be a finitely generated ideal such that A/I is finitely generated as \mathbb{Z} -module. Assume that there exists a constant d such that $\dim_F(A \otimes F) = \dim_F(A/I \otimes F) = d$ for any field F . Then $I = 0$ and A is a free \mathbb{Z} -module of rank d .

Proof. If M is a finitely generated \mathbb{Z} -module such that $\dim_F(M \otimes F) = d$ for all fields F , then it follows from the structure theorem for finitely generated modules over a principal ideal domain that M is a free \mathbb{Z} -module of rank d . In particular, A/I is a free \mathbb{Z} -module of rank d by our assumption, and it suffices to show that $I = 0$.

As \mathcal{Q} is \mathbb{Z} -flat, we have an exact sequence

$$0 \rightarrow I \otimes \mathcal{Q} \rightarrow A \otimes \mathcal{Q} \rightarrow A/I \otimes \mathcal{Q} \rightarrow 0.$$

Since $A \otimes \mathcal{Q}$ and $A/I \otimes \mathcal{Q}$ are of the same finite dimension over \mathcal{Q} by assumption, we have $I \otimes \mathcal{Q} = 0$. This implies that I is a torsion group. Since I is finitely generated as ideal, there exists an integer $m > 0$ such that $mI = 0$.

We now fix a prime p , and let $\mathbb{Z}_{(p)}$ be the localization of \mathbb{Z} at $p\mathbb{Z}$ and \mathbb{Z}_p its residue field. Then $\mathbb{Z}_{(p)}$ is flat over \mathbb{Z} and there exists an exact sequence

$$0 \rightarrow \mathbb{Z}_{(p)} \xrightarrow{p \cdot} \mathbb{Z}_{(p)} \rightarrow \mathbb{Z}_p \rightarrow 0,$$

where $p \cdot$ denotes multiplication. Thus we obtain the following exact commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & I \otimes \mathbb{Z}_{(p)} & \longrightarrow & A \otimes \mathbb{Z}_{(p)} & \longrightarrow & A/I \otimes \mathbb{Z}_{(p)} \longrightarrow 0 \\ & & p \cdot \downarrow & & p \cdot \downarrow & & p \cdot \downarrow \\ 0 & \longrightarrow & I \otimes \mathbb{Z}_p & \longrightarrow & A \otimes \mathbb{Z}_p & \longrightarrow & A/I \otimes \mathbb{Z}_p \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & I \otimes \mathbb{Z}_p & \longrightarrow & A \otimes \mathbb{Z}_p & \longrightarrow & A/I \otimes \mathbb{Z}_p \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0. \end{array}$$

As A/I is free over \mathbb{Z} , the $\mathbb{Z}_{(p)}$ -module $A/I \otimes \mathbb{Z}_{(p)}$ is free as well. Hence $p \cdot : A/I \otimes \mathbb{Z}_{(p)} \rightarrow A/I \otimes \mathbb{Z}_{(p)}$ is a monomorphism. So the Snake Lemma yields an exact sequence

$$0 \rightarrow I \otimes \mathbb{Z}_p \rightarrow A \otimes \mathbb{Z}_p \rightarrow A/I \otimes \mathbb{Z}_p \rightarrow 0.$$

Since $A \otimes \mathbb{Z}_p$ and $A/I \otimes \mathbb{Z}_p$ are of the same finite dimension over \mathbb{Z}_p , we have $I \otimes \mathbb{Z}_p = 0$. On the other hand, we have $I \otimes \mathbb{Z}_p \cong I/pI$. Hence $pI = I$, for any prime p . Therefore, $I = mI = 0$, and the proof of the lemma is complete.

REMARK. The above lemma fails for the trivial extension $\mathbb{Z} \ltimes (\mathbb{Q}/\mathbb{Z})$, whence finite generation of the ideal I is necessary.

2.4. PROPOSITION. *Let R be a commutative ring and Γ a finite Auslander-Reiten quiver. Then the mesh algebra $R(\Gamma)$ is a free R -module of finite rank and that rank is independent of R .*

Proof. In this proof, all tensor products are taken over \mathbb{Z} . Note first that $R(\Gamma) \cong \mathbb{Z}(\Gamma) \otimes R$. It is thus sufficient to show that $\mathbb{Z}(\Gamma)$ is free over \mathbb{Z} of finite rank.

We first fix an algebraically closed field k . Since $k(\Gamma)$ is finite dimensional over k , there exists $r > 0$ such that $\mathfrak{R}^r k(\Gamma) = 0$. Set $A = \mathbb{Z}(\Gamma)$ and $I = \mathfrak{R}^r \mathbb{Z}(\Gamma)$. Then I is finitely generated as an ideal and A/I is finitely generated as \mathbb{Z} -module. Let F be an arbitrary field. Under the isomorphism $A \otimes F \cong F(\Gamma)$, the image of $I \otimes F$ in $A \otimes F$ maps onto $\mathfrak{R}^r F(\Gamma)$. Hence, $A/I \otimes F \cong F(\Gamma)/\mathfrak{R}^r F(\Gamma)$. On the other hand, by Theorem 2.2, $\dim_F \mathfrak{R}^n F(\Gamma) = \dim_k \mathfrak{R}^n k(\Gamma)$ for all $n \geq 0$. In particular, $\mathfrak{R}^r F(\Gamma) = 0$. This yields

$$\dim_F (A/I \otimes F) = \dim_F F(\Gamma) = \dim_F (A \otimes F),$$

which is further equal to $\dim_k k(\Gamma)$. Now Lemma 2.3 applies and shows that $\mathbb{Z}(\Gamma) = A$ is a free \mathbb{Z} -module of finite rank. This completes the proof.

Acknowledgements. Both authors gratefully acknowledge partial support from the Natural Sciences and Engineering Research Council of Canada.

References

- [1] M. AUSLANDER, I. REITEN, and S. O. SMALØ, “Representation Theory of Artin Algebras”, Cambridge Studies in Advanced Mathematics **36** (Cambridge University Press, Cambridge 1995).
- [2] R. BAUTISTA, P. GABRIEL, A. V. ROITER, and L. SALMERÓN, “Representation-finite algebras and multiplicative basis”, *Invent. Math.* **81** (1985) 217 - 285.
- [3] K. BONGARTZ and P. GABRIEL, “Covering spaces in representation theory”, *Invent. Math.* **65** (1982) 331 - 378.

- [4] O. BRETSCHER and P. GABRIEL, “The standard form of a representation-finite algebra”, *Bull. Soc. Math. France* **111** (1983) 21 - 40.
- [5] S. BRENNER, “A combinatorial characterisation of finite Auslander-Reiten quivers”, *Lecture Notes in Mathematics* **1177** (Springer-Verlag, Berlin 1986) 13 - 49.
- [6] R.-O. BUCHWEITZ and S. LIU, “Hochschild cohomology and representation-finite algebras”, preprint.

Ragnar-Olaf Buchweitz
Department of Mathematics
University of Toronto
Toronto, Ontario
Canada M5S 3G3
ragnar@math.toronto.edu

Shiping Liu
Département de mathématiques
Université de Sherbrooke
Sherbrooke, Québec
Canada J1K 2R1
shiping@dmi.usherb.ca